

## Indi Mid-Maturing MIN-DAK Region



## **NAVY BEAN**


**Indi** – Mid-Maturing

## **Profile:**

Indi is a mid-maturing, indeterminate upright vine which is adapted to MN/DAK. Indi's upright structure gives good pod clearance and makes it a variety that is highly suitable for direct harvest. The Indi variety has consistently shown erect structure, improved pod clearance, resistance to lodging, and excellent concurrent dry-down of pods.

## **Economic Profile:**

Assuming a 287 lbs/AC yield advantage compared to Ensign, profit would be \$91.84/AC on \$32.00/cwt beans minus the difference in seed cost/AC. This can calculate out to an approximate \$14,694 increased profit on one quarter section field.



<sup>\*</sup>Indi is a licensed ADM Edible Bean Specialties, Inc. variety (PVP).



TRIAL	DATA	
INDI		
Approx. Maturity	96-101 days	
Plant Type	2A	
Approx. Seed Count	2,494 seeds/lb.	
Disease Resistance	IR:BCMV ("I" gene),	
	Ua	

See reverse side for disease resistance abbreviation chart

For customers around the world, ADM draws on its resources—its people, products, and market perspective—to help them meet today's consumer demands and envision tomorrow's needs.



<sup>\*\*</sup> Ensign is a UC-Davis, public variety.





KEY TO RESISTANCE ABBREVIATIONS FOR BEANS	
Plant Type 1A	Bush determinate erect stem
Plant Type 2A	Erect growth indeterminate short runners
Plant Type 2B	Erect growth indeterminate with medium to long runners
Plant Type 3B	Prostrate vine indeterminate growth with long runners
BCMV	Bean common mosaic caused by the specified strains of Bean common mosaic virus
BCTV	Curly top caused by Beet curly top virus
всуму	Bean golden yellow mosaic caused by Bean golden yellow mosaic virus
CI	Anthracnose caused by Collectrichum lindemuthianum
Psp	Halo blight caused by Pseudomnas savastanoi pv. phaseolicola
Pss	Bacterial brown spot caused by Pseudomaonas syringae pv. syringae
Ua	Rust caused by the specified races of Uromyces appendiculatus
HR	<b>High Resistance:</b> describes plant varieties that highly restrict the growth and development of the specified pest or pathogen under normal pest or pathogen pressure when compared to susceptible varieties. Highly resistant varieties may, however, exhibit some symptoms or damage under heavy pest or pathogen pressure.
IR	Intermediate Resistance: describes plant varieties that restrict the growth and development of the specified pest or pathogen, but may exhibit a greater range of symptoms or damage compared to highly resistant varieties. Intermediately resistant varieties will still show less severe symptoms or damage than susceptible plant varieties when grown under similar environmental conditions and/or pest or pathogen pressure.

In cases where specific races or strains are not noted the variety is resistant to some, but not necessarily all known races or strains of the pathogen.

**ADM Seedwest** 

P.O. Box 1470

Decatur, Illinois 62525

ADM