Pinto 'Torreon'




Direct harvestable high yielding upright pinto bean

PROFILE

Torreon has shown broad adaptability in trials throughout the High Plains, Idaho and in the MN-DAK region. Torreon is indeterminate, upright in its architecture providing great clean quality product and provides great yield potential to the grower.

Torreon has an excellent upright architecture for direct harvest. Torreon also has excellent seed color, appearance and shape.

Torreon has been yielding competitively with Monterrey while maturing out 1 day earlier, and has good dry down characteristics, aiding in its harvestability.



- * Yield data based on 37 yr./locations of data.
- ** Maturity data is based on 23 yr./locations of data.
- *** Seed count data is based on 37 yr./locations of data.

 Data from 2007 2016 trials in Buxton, Forest River, Hatton, Prosper,
 Northwood. St. Thomas and more.



TRIAL DATA TORREON

Approx. Maturity Matures approx. 1 day earlier than Monterrey**

Approx. Seed Count 1,213 seeds/lb.***

Disease Resistance Resistance Resistant to Rust strains tested for at Colorado State University

TO PURCHASE SEED: CONTACT YOUR LOCAL DEALER

For customers around the world, ADM draws on its resources—its people, products, and market perspective—to help them meet today's consumer demands and envision tomorrow's needs.



Pinto 'Torreon'



Direct harvestable high yielding upright pinto bean

PT Torreon was developed through ProVita, Inc.'s pinto breeding program through a contract with ADM—Seedwest and is the sole property of ADM—Seedwest.

> Plant Variety Protection for PT Torreon is applied for. Unauthorized propagation of this variety is prohibited.

All variety information presented herein is based on field and laboratory observations. Actual crop yield and quality are dependent upon many factors beyond our control. Since environmental conditions and local practices may affect variety characteristics and performance, we disclaim legal responsibility therefore. Read all tags and labels. They contain important conditions of sale, including limitations of warranties and remedies.

KEY TO RESISTANCE ABBREVIATIONS FOR BEANS

Plant Type 1A	Bush determinate erect stem
Plant Type 2A	Erect growth indeterminate short runners
Plant Type 2B	Erect growth indeterminate with medium to long runners
Plant Type 3B	Prostrate vine indeterminate growth with long runners
BCMV	Bean common mosaic caused by the specified strains of Bean common mosaic virus
BCTV	Curly top caused by Beet curly top virus
BGYMV	Bean golden yellow mosaic caused by Bean golden yellow mosaic virus
CI	Anthracnose caused by Collectrichum lindemuthianum
Psp	Halo blight caused by Pseudomnas savastanoi pv. phaseolicola
Pss	Bacterial brown spot caused by Pseudomaonas syringae pv. syringae
Ua	Rust caused by the specified races of Uromyces appendiculatus
HR	High Resistance: describes plant varieties that highly restrict the growth and development of the specified pest or pathogen under normal pest or pathogen pressure when compared to susceptible varieties. Highly resistant varieties may, however, exhibit some symptoms or damage under heavy pest or pathogen pressure.
IR	Intermediate Resistance: describes plant varieties that restrict the growth and development of the specified pest or pathogen, but may exhibit a greater range of symptoms or damage compared to highly resistant varieties. Intermediately resistant varieties will still show less severe symptoms or damage than susceptible plant varieties when grown under similar environmental conditions and/or pest or pathogen pressure.

In cases where specific races or strains are not noted the variety is resistant to some, but not necessarily all known races or strains of the pathogen.

Rev. 09-10-2021

